

	Drug ¹			Placebo ²			Comments
	Dropout 6 weeks	Rx Lost	10 lb Range loss (pounds)	Dropout 6 weeks	Rx Lost	10 lb Range (pounds)	
2—Gremmel No. 2537:							
High.....	3	7	3	3	8	0	11-39 dropouts: Relatively few lapses in therapy or intercurrent illnesses report.
Low.....	2	6	1	3	7	0	
Total.....	5	13	4	6	15	0	
3—Esposito No. 2539:							
High.....	3	6	0	1	7	0	4-33 dropouts: Irregular treatment administration during holidays and intercurrent flu.
Low.....	0	8	1	0	8	0	
Total.....	3	14	1	1	15	0	
4—Steri No. 2540:							
High.....	1	7	0	1	8	0	5-31 dropouts: High incidence of intercurrent illness. Frequent lapses of treatment administration. Protocol deviation indicated by investigator on 4 case records.
Low.....	1	7	1	2	4	0	
Total.....	2	14	1	3	12	0	
5—Gottreau No. 2543:							
High.....	2	6	1	3	2	0	12-29 dropouts: Only 17 patients evaluated. Many subjects had intercurrent illness.
Low.....	2	4	0	5	5	0	
Total.....	4	10	1	8	7	0	
6—Noble No. 2: No. 2545:							
High.....	1	7	4	2	6	1	8-32 dropouts: This investigator is credited with 2 studies conducted simultaneously. However, No. 2544 cannot be evaluated due to illegible records, irregular monitoring, confused identity of which treatment patient received. No. 2545 demonstrated greatest efficacy reported for both drug and placebo.
Low.....	3	5	2	2	6	1	
Total.....	4	12	6	4	12	2	12, 18
Grand total.....	18	63	13	22	61	2	12, 18

¹ 81 subjects initially.

² 83 subjects initially.

³ 12 lost 11-14½ lb; 1 lost 31 lb.