

Promotion

8. Parke Davis Chloromycetin entries in Vademecum Daimon, Nov. 1972.
Spanish physicians' desk reference published by pharmaceutical companies.

Spain, Barcelona November 1972
Parke-Davis entries in Vademecum Daimon (Spain) PD

CHLOROMYCETIN

Composition: Chloramphenicol Parke Davis.

(physicians desk reference published by pharmaceutical companies)

ACTION: Antibiotic of ample spectrum, which covers a large number of bacteria--gram negative and gram-positive-- all the microorganisms of the rickettsia, lymphogranuloma and inguinal as well as psittacosis granuloma.

Indications: Typhoid fever and other salmonellas, bacillar dysentery rickettsiosis, infections of the urinary tract caused by a variety of gram negative and gram positive bacteria, serious surgical infections, including peritonitis, some respiratory infections, meningitis, venereal diseases such as pharyngitis resistant to other antibiotics, inguinal granuloma and venereal lymphogranuloma. Septicemias, brucellosis, psittacosis, etc.

Administration: 50 mg/kg/day, in 4 fractionated doses every 6 hours. For premature and newly born babies and others with physiological immaturity, 25/mg/kg/day.

CAUTION: If it is possible, it is convenient to have hemograms before establishing the Chloromycetin therapy and they should be repeated if a prolonged or intermittent administration is necessary, or in cases when the dose given is higher than usual, in order to prevent a possible depression of the bone marrow. Although rarely, some irreversible discrasias have been observed. The "gray syndrome" has been described in premature and newly born babies.

As it concerns to hypersensitive reactions, some febrile reactions, cutaneous eruptions and allergic reactions can be observed.

For more information on adverse reactions and indications please write to the laboratory for a monograph of Chloromycetin.

***Presentation:**

CLOROSTREP (capsuls and liquid)

Composition: Each capsule or tea spoon contains 125 mg Of Chloromycetin and 125 mg of dihydrostreptomycin.

Indications: Salmonella and shigella infections, ulcerous colitis, gastroenteritis, pre and post operatory intestinal tract, etc.

Dose: Children: 1 or 2 tea spoons every 6 hr, adults 2-4 capsules every 6 hours.

Presentation: 8 Capsules in a container. Liquid in a 60 cc bottle.

***Presentation:** Capsules of 250 mg, 12 in a container; 122,60 pts with 24; 228.90 pts. Cream at 1%, tube with 30 g, 79.20 pts. Topical at 10%, 5 cc container, 98.40 pts. Intramuscular: vials with 1 g, 96.40 pts; with 2 g. 166 pts. Succinate: vial 1 g, 86.30 pts.