

COMPETITIVE PROBLEMS IN THE DRUG INDUSTRY 15521

Chloromycetin

Dose:

In general a dose of 50 mg/kg/day is recommended, in 4 doses given with intervals of 6 hours. In serious cases like septicemia and meningitis the dose could be increased to 100 mg/kg/day.

For premature babies and babies up to a month old it is recommended not to give over 25 mg/kg/day due to their physiological immaturity.

Chloromycetin - flask with 12 capsules P.V.P.	122.60	(pesetas)
" " " 24 "	228.90	"
Palmitate of Chloromycetin: flask with 60 cc	129.90	"
Topical chloromycetin: Flask with 5 cc	76.10	pesetas

For more information about prescribing the product, request the monograph which will be available to you and which describes indications and adverse reactions of the product.

Spain, Barcelona November 1972
Parkelase tubes in Vademecum Dainion, Spain's PDR

Parkelase with Chloromycetin

Composition: Combination of two ^{lytic} enzymes, fibrinolisin and desoxyrrribo-nuclease with chloramphenicol, in ointment.

Action: Enzymatic and antibiotic debridement.

Indications: The same as for Parkelase when an infection - a complication - is suspected or exists. Burns, wounds, cervicits, vaginitis, ulcers etc.

Dose: 1 to 3 daily applications.

Presentation: Ointment - tubes with 10 and 30 grams.

1-Dosification - "See attached prospect (literature)"

2. "Algunas" some respiratory infections

3. "Comoblennoragia"