

15542 COMPETITIVE PROBLEMS IN THE DRUG INDUSTRY

Acquired in:
Spain, Barcelona June 1974
(translation of glossy colored picture ad)

CHLOROSTREP - Chloramphenicol and Sulfate of Dihydrostreptomycin IT ATTACKS BY TWO FRONTS

GENERAL
Chloramphenicol

LOCAL
Dihydrostreptomycin
and chloramphenicol

When chloramphenicol is administered by the oral route, the major part is absorbed in the intestine, it distributes itself among the tissues and liquids in the organism and exercises an antibiotic action in general; the amount not absorbed, however, is enough to give a concentration therapeutically acceptable in the intestine. When streptomycin is administered by the oral route, practically it is not absorbed in the intestine and reaches it in a very high concentration.

DOSE

Children: 1 to 2 teaspoonfuls every 6 hours
Adults: 2 capsules every 4 - 6 hours

PRESENTATION:

Bottle of 8 capsules P.V.P. 119.80
Bottle of 60 cc. P.V.D. 160/40

(Back page)

CHLOROSTREP - Chloramphenicol and Sulfate of Dihydrostreptomycin

WARNING

It is recommended to use Chlorostrep only under the direction of a physician. It is not recommended to use it in therapy of trivial (minor) infectious processes. For complete prescribing information regarding chloramphenicol, including warnings, precautions and adverse reactions - amongst them depression of the bone marrow - please see the monograph on chloramphenicol available on request.

WITH THE GUARANTEE OF

PARKE-DAVIS
MADRID- BARCELONA

CHLST-V-74 (D.M.I) Grafiberica Marques de Cadiz 25 Neres Dep Legal 27- 1974