

*Spain, Barcelona May, 1973
Translation of article, *Journal of the American Medical Association*, published.*

CHLOROSTREP - Chloramphenicol and Sulfate of Dihydroestrptomycin.

Capsules P.V.P. 119,80 LOCAL DIHYDROESTRPTOMYCINE

Liquid P.V.P. 159,60 GENERAL Chloromycetin

It Broadens the Bacteria Spectrum for a Greater Clinic Efficacy

The association of chloramphenicol and streptomycin has a broader bacteria spectrum than any other antibacterian agent clinically used alone. This spectrum extremely broad, attacks the microorganisms that commonly cause the infectious diarrhea, enteric and mix infections that can occur in the postoperative surgery

It attacks by two fronts:

When chloramphenicol is administered by the oral route, the majority is absorbed in the intestine, it distributes itself among the tissues and liquids in the organism and exercises an antibiotic action in general; the amount not absorbed, however, is enough to give a concentration therapeutically accepted in the intestine. When (strept) is administered by the oral route, practically is not absorbed in the intestine and reaches in it a very high concentration.

For complete information on prescriptions including indications and adverse reactions, write for the monograph.

MADRID, BARCELONA

Advertising page: Summer and Infectious Diarrheas:

Your vacation can be more complete with

CHLOROSTREP

It attacks by two fronts. PARKE DAVIS