

COMPETITIVE PROBLEMS IN THE DRUG INDUSTRY 16605

82. Salguero, C.H., et al., "Propoxyphene Dependence," JAMA 210: 135, 1969.
83. Drug Control Division, supra note 24, pp 61-62.
84. Klein, R.B., et al., "Probable Neonatal Propoxyphene Withdrawal: A Case Report," Pediatrics 55(6):882, 1975.
85. Quillian, W.W., and C.A. Dunn, "Neonatal Drug Withdrawal from Propoxyphene," JAMA 235: 2128, 1976.
86. Tyson, H.K., "Neonatal Withdrawal Symptoms Associated With Maternal Use of Propoxyphene Hydrochloride," J Pediatrics 85: 684, 1974.
87. Maletsky, B.M., "Addiction to Propoxyphene (Darvon): A Second Look," Int J Addict 9(6): 775, 1974.
88. Maletsky, B.M., supra note 87; Tennant, F.S., supra notes 72 and 73.
89. Tennant, F.S., supra note 15.
90. Chambers, C.D., et al., "Five Patterns of Darvon Abuse," Int J Addict 6(1): 173, 1971.
91. Federal Register 42(29): 8636, February 11, 1977.
92. Meyers, F.H., E. Jawetz, and A. Golfien, Review of Medical Pharmacology, 5th ed., Lange Medical Publications, Los Altos, California, 1976, p 295.
93. Goodman, L., and A. Gilman, The Pharmacologic Basis of Therapeutics, 5th ed., MacMillan Publishing Co., Inc., New York, 1975, p.305.
94. Miller, R.R., et al., "Propoxyphene Hydrochloride: A Critical Review," JAMA 213: 996, 1972.
95. Physicians' Desk Reference, 32nd ed., 1978, p.998.
96. Moertel, C.G., et al., "A Comparative Evaluation of Marketed Analgesic Drugs," N Eng J Med 286: 813, 1972.
97. Hopkinson, J.H., III, et al., "Acetaminophen Versus Propoxyphene Hydrochloride For Relief of Pain in Episiotomy Patients," J Clin Pharmacol 13:251, 1973.
98. Gruber, C.M., "Codeine and Propoxyphene in Postepisiotomy Pain," JAMA 237: 2734, 1977.
99. Moertel, C.G., et al., supra note 96.
100. Gruber, C.M., "Effectiveness of Propoxyphene," (letter) N Eng J Med 286: 1158, 1972.
101. Gruber, C.M., et al., "Comparative Evaluation of Analgesic Agents in Postpartum Patients: Oral Dextropropoxyphene, Codeine and Meperidine," Anesth Analg 41: 538, 1962; see also supra note 93.