3. Thiazide compounds should be administered with caution to patients who

show elevated Urea Nitrogen (BUN).

4. Patients on tranquilizers may experience faintness due to the development of orthostatic hypotension. The blood pressure of these patients should be checked in a sitting or standing position. If low, the tranquilizers should be discontinued, or the patient kept in bed.

The following is an addendum (as of April 1, 1967) of all additions or dele-

tions to the Formulary since October 1, 1966.

New drugs

1. Auralgan Ear Drops 15 ml.

Chlordantoin Cream (Sporostacin)

3. Ethacrynic Acid (Edecrin), Tablets-Capsule shaped, 25 mg., 50 mg.; Injection-for I.v. use 50 mg. vial.

4. Fibrinolysin and Desoxyribonucleas Combined-Bovine (Elase Ointment) 30 Gm. tubes

- 5. Furosemide (Lasix) 40 mg. tablets
- 6. Hyperphos-oral

7. Imphos-intravenous

- 8. Medroxyprogesterone Acetate (Depo-Provera) Inj.
- 9. Polymyxin B (Aerosporin) Otic Solution 10 ml.
- 10. Pseudoephedrine HCl (Sudafed) Syrup, 10 ml.

New dosage forms

- 1. Amitriptyline HCl (Elavil) 50 mg. tablets.
- 2. Ampicillin: injection, 500 mg. vial; Capsule, 500 mg.; Suspension, 250 mg./
- 3. Atropine Sulfate Ophthamlic Ointment 1%.
- 4. Bethanechol Chloride 25 mg. tab-
- 5. Chloroprocaine HCL (Nesacaine)
- 6. Dexamethasone Elixir 0.5 mg./ 5 ml.
- 7. Echothiophate (Phospholine Iodide) 0.25%.

- 8. Methenamine Mandelate 1 tablets.
- 9. Methenamine Suspension 500 mg./ 5 ml. Note: Replaces the 250 mg./5 ml.
- 10. Methotrexate Sodium Inj. 5 mg.
- 11. Penicillin G. Potassium 250 mg. tablets.
- 12. Penicillin, Potassium Phenoxymethyl-250 mg. tablets.
- 13. Perphenazine (Trilafon) 16 mg. tablets.
- 14. Trifluoperazine (Stelazine) 10 mg.

Drugs available for special departments only

- 1. Sparteine Sulfate Inj. 150 mg. ampul; OB-GYN Service.
- 2. Isoxsuprine (Vasodilan), Injection 5-mg./ml., 2 ml. ampul; OB-GYN Serv-
- 3. X-Prep, Radiology Service of requesting hospitals only.
- 4. Hydroxyzine (Vistaril), tion-25 mg. disposable syringe; Neurosurgery Service.

Deletions

- 1. Streptokinase Streptodornase (Varidase) Jelly. Note: Replaced by Elase Ointment.
- 2. Ampicillin Suspension, 125 mg./5 ml. Note: Replaced by 250 gm./5 ml.
- 3. Methenamine Suspension 250 mg. 5 ml. Note: Replaced by 500 mg./5 ml.

ar rayed **al**e

Please cut along dotted line, fold in half and place in front of your Formulary. Do not discard previous addendum.

New pages for the Formulary replacing the addendum will be available by the end of April.

Pharmacy Bulletin-Published quarterly as a service to the Medical Staff of Montefiore Hospital and Medical Center by the Pharmacy Department.

Appreciation is expressed to Dr. Francis Foldes and Dr. Milton Reisch for their assistance in the preparation of this issue of the Pharmacy Bulletin.

Kurt Kleinmann, M.Sc., Director of Pharmacy Service.

Robert J. Petrick, M.Sc., Asst. Director. Margaret S. Oppedal, M.Sc., Supervisor. Louis Kalaboke, B.S., Supervisor.

Zina Fediay, M.Sc., Supervisor.

Senator Nelson. Our next witness is Dr. Calvin M. Kunin, associate professor of preventive medicine and medicine, University of Virginia School of Medicine. Doctor, your statement will be placed in full in the record, including your professional credentials. You may proceed to summarize your statement and, if you don't mind, I may occasionally