

[From Antibiotics and Chemotherapy magazine, Nov. 1954]

when resistance to other antibiotics develops...

Chloromycetin®

Current reports^{1,2} describe the increasing incidence of resistance among many pathogenic strains of microorganisms to some of the antibiotics commonly in use. Because this phenomenon is often less marked following administration of CHLOROMYCETIN (chloramphenicol, Parke-Davis), this notably effective, broad spectrum antibiotic is frequently effective where other antibiotics fail.

Coliform bacilli—100 strains

up to 43% resistant to other antibiotics;
2% resistant to CHLOROMYCETIN.¹

Staphylococcus aureus—500 strains

up to 73% resistant to other antibiotics;
2.4% resistant to CHLOROMYCETIN.²

CHLOROMYCETIN is a potent therapeutic agent and, because certain blood dyscrasias have been associated with its administration, it should not be used indiscriminately or for minor infections. Furthermore, as with certain other drugs, adequate blood studies should be made when the patient requires prolonged or intermittent therapy.

References

(1) Kirby, W. M. M.; Waddington, W. S., & Doornink, G. M.: Antibiotics Annual, 1953-1954, New York, Medical Encyclopedia, Inc., 1953, p. 285. (2) Finland, M., & Haight, T. H.: *Arch. Int. Med.* 91: 143, 1953.

PARKE, DAVIS & COMPANY



DETROIT 32, MICHIGAN