


[From General Practitioner magazine, May 1961]



IN VITRO SENSITIVITY OF 250 STRAINS OF STAPHYLOCOCCI
TO CHLOROMYCETIN AND TO FOUR OTHER ANTIBIOTICS*

	CHLOROMYCETIN 78%
	Antibiotic A 68%
	Antibiotic B 55%
	Antibiotic C 45%
	Antibiotic D 21%

These strains of coagulase-positive staphylococci were isolated from hospitalized patients at a large county hospital during the year 1959. Sensitivity tests were done by the disc method.

*Adapted from Bauer, Perry, & Kirby¹

References: (1) Bauer, A. W.; Perry, D. M., & Kirby, W. M. M.: *J. A. M. A.* 173:475, 1960. (2) Fisher, M. W.: *Arch. Int. Med.* 105:413, 1960. (3) Cohen, S.: *Circulation* 20:96, 1959. (4) Edwards, T. S.: *Am. J. Ophth.* 48, Part II:19, 1959. (5) Smith, I. M.: *Staphylococcal Infections*, Chicago, The Year Book Publishers, Inc., 1958, p. 148. (6) Petersdorf, R. G.; Rose, M. C.; Minchew, H. B.; Keene, W. R., & Bennett, I. L., Jr.: *Arch. Int. Med.* 105:398, 1960. (7) Editorial: *J. A. M. A.* 173:544, 1960. (8) Finland, M.; Jones, W. E., Jr., & Bennett, I. L., Jr.: *Arch. Int. Med.* 101:363, 1959.

PARKE-DAVIS

PARKE, DAVIS & COMPANY, Detroit 22, Michigan