

TABLE 6.—REPORTS RECEIVED OF LEUKEMIA FOLLOWING CHLORAMPHENICOL OR PHENYLBUTAZONE THERAPY—SUMMARY OF CASES NOT PREVIOUSLY REGISTERED FOR ADVERSE REACTION¹

Case:	Sex		Leukemia		Drug ingestion			Initial reaction		Latent period
	Age, years	Type	Type	Latent period	Drug	Indication	Duration	Estimated dose, grams	Type	
1. 60 F	60	AML ²	2 years	12-15 years (10 months after initial reaction)	C	Frequent common colds	Intermittently—14 months	15	Leukopenia	23 months
2. 56 F	56	AML ²	12-15 years (10 months after initial reaction)		C	Cystitis	Intermittently—12-15 years	200	Erythroid hypoplasia, leukopenia (marrow-maturation arrest of WBC)	12-15 years
3. 64 F	64	CML	9½ years	12 years (7 months after initial reaction)	C	Sinusitis	1 week	7	Pancytopenia	1 month
4. 61 F	61	SML	12 years		P	Rheumatoid arthritis	Intermittently—11-12 years	21	Leukopenia	11-12 years
5. 43 F	43	AML ²	1 year		P	Bursitis	Intermittently—9 months	> 15	Leukopenia (marrow-maturation arrest of WBC)	9 months
6. 77 F	77	AML ²	3½ years		P	Rheumatoid arthritis	Intermittently—3 years	> 50	Leukopenia	3 years

¹ See table 5 footnote for explanation of symbols. In addition, SML = subacute myelogenous leukemia. ² Termed "aleukemic" leukemia at diagnosis.