

Precautions: ANTIHYPERTENSIVE THERAPY WITH HYGROTON SHOULD ALWAYS BE INITIATED CAUTIOUSLY in postsympathectomy patients and IN PATIENTS RECEIVING GANGLIONIC BLOCKING AGENTS OR OTHER POTENT ANTIHYPERTENSIVE DRUGS, or curare. Reduce dosage of concomitant antihypertensive agents by at least one-half. Barbiturates, narcotics or alcohol may potentiate hypotension. BECAUSE OF THE POSSIBILITY OF PROGRESSION OF RENAL DAMAGE, PERIODIC DETERMINATION OF THE BUN IS INDICATED. Discontinue if the BUN rises or liver dysfunction is aggravated. HEPATIC COMA MAY PRECIPITATED.

Electrolyte imbalance, SODIUM AND/OR potassium depletion may occur. IF POTASSIUM DEPLETION SHOULD OCCUR DURING THERAPY, HYGROTON SHOULD BE DISCONTINUED AND POTASSIUM SUPPLEMENTS GIVEN, PROVIDED THE PATIENT DOES NOT HAVE MARKED OLIGURIA.

Take special care in cirrhosis or severe ischemic heart disease and in patients receiving corticosteroids, ACTH, or digitalis. Salt restriction is not recommended.

Adverse Reactions: Nausea, gastric irritation, vomiting, anorexia, constipation and cramping, dizziness, weakness, restlessness, hyperglycemia, hyperuricemia, headache, muscle cramps, orthostatic hypotension, aplastic anemia, leukopenia, thrombocytopenia, agranulocytosis, impotence, dysuria, transient myopia, skin rashes, urticaria, purpura, necrotizing angitis, ACUTE GOUT, AND PANCREATITIS WHEN epigastric pain or UNEXPLAINED G.I. symptoms DEVELOP after prolonged administration. Other reactions reported with this class of compounds include: jaundice, xanthopsia, paresthesia, and photosensitization.

Average Dosage: One tablet (100 mg.) with breakfast daily or every other day.

Availability: White, single-scored tablets of 100 mg. in bottles of 100 and 1000.

(NOTE.—This revised "Brief Summary", for use in future medical journal advertising, contains additional words and phrases (printed in capital letters) taken from the official package insert.)

BRIEF SUMMARY OF REGROTON®—CHLORTHALIDONE, 50 MG., RESERPINE U.S.P., 0.25 MG.

Indications: Hypertension.

Contraindications: History of mental depression, hypersensitivity, and most cases of severe renal or hepatic diseases.

Warning: With the administration of enteric-coated potassium supplements, WHICH SHOULD BE USED ONLY WHEN ADEQUATE DIETARY SUPPLEMENTATION IS NOT PRACTICAL, the possibility of small bowel lesions (OBSTRUCTION, HEMORRHAGE, AND PERFORATION) should be kept in mind. SURGERY, FOR THESE LESIONS HAS FREQUENTLY BEEN REQUIRED AND DEATHS HAVE OCCURRED.

DISCONTINUE COATED POTASSIUM-CONTAINING FORMULATIONS IMMEDIATELY IF ABDOMINAL PAIN, DISTENTION, NAUSEA, VOMITING, OR GASTROINTESTINAL BLEEDING OCCUR.

Use cautiously during pregnancy since adverse reactions (thrombocytopenia, hyperbilirubinemia, altered carbohydrate metabolism, etc.) are potential problems in the newborn.

Discontinue 2 weeks before general anesthesia, 1 week before electroshock therapy, and if depression or peptic ulcer occurs.

Precautions: ANTIHYPERTENSIVE THERAPY WITH REGROTON SHOULD ALWAYS BE INITIATED CAUTIOUSLY in postsympathectomy patients and IN PATIENTS RECEIVING GANGLIONIC BLOCKING AGENTS, OTHER POTENT ANTIHYPERTENSIVE DRUGS, or curare. Reduce dosage of concomitant antihypertensive agents by at least one-half.

BECAUSE OF THE POSSIBILITY OF PROGRESSION OF RENAL DAMAGE, PERIODIC KIDNEY FUNCTION TESTS ARE INDICATED. Discontinue if the BUN rises or liver dysfunction is aggravated, HEPATIC COMA MAY BE PRECIPITATED.

Electrolyte imbalance, SODIUM AND/OR potassium depletion may occur. IF POTASSIUM DEPLETION SHOULD OCCUR DURING THERAPY, REGROTON SHOULD BE DISCONTINUED AND POTASSIUM SUPPLEMENTS GIVEN, PROVIDED THE PATIENT DOES NOT HAVE MARKED OLIGURIA.

Take particular care in cirrhosis or severe ischemic heart disease and in patients receiving corticosteroids, ACTH, or digitalis. Salt restriction is not recommended. BILIARY COLIC MAY BE PRECIPITATED (IN PATIENTS