of the questionnaire, more than 11,800 responses had been received. Of these, 11,603 were satisfactory for data processing and tabulation: 3,200 from general practitioners, 1,784 from internists, 1,005 from surgeons, 710 from pediatricians, 380 from orthopedic surgeons, 70 from physiatrists, and 4,454 from physicians of other specialties (Table 4).

Almost all responders, classified according to their specialty or primary type of practice, are represented in about the same proportion in the survey as in the national distribution of the physician population (Table 4). Among the general practitioners, 89.1% reported treating patients with rheumatoid arthritis, among the internists 90%, among the physiatrists 90%, and among the orthopedic surgeons 82.9%.

Table 5. Regional distribution of physicians replying to questionnaire and of all physicians in the United States

Region	Physicians replying to questionnaire		All physicians in U.S.*
	Number	Percent	Percent
New England	722	6.2	6.6
Mid-East	2,291	19.7	23.0
South Atlantic	1,413	12.2	12.7
Great Lakes	1,816	15.7	18.0
Mid-South	481	4.1	4.8
Plains	955	8.2	7.2
Southwest	856	7.4	7.9
Rocky Mountain	549	4.7	3.8
Far West	1,793	15.5	15.5
Pacific	55	0.5	0.5
Unknown	672	5.8	
Total *Source: Buckley Dement	11,603	100.0	100.0