

CRITERIA FOR EVALUATING CLINICAL RESULTS

UPPER AND LOWER RESPIRATORY INFECTIONS

*Soft-tissue Infections**Miscellaneous infections*

Favorable responses: Includes those designated as "good" or "satisfactory."

Good—patient showed definite favorable response to doxycycline therapy with prompt alleviation of symptoms.

Satisfactory—patient showed beneficial response, but the duration of symptoms was longer than might have been expected with a good response.

Poor responses: Cases in which it was felt there was no response, or a worsening of symptoms.

Infections of the genitourinary tract

Favorable responses: Includes those designated as "good" or "satisfactory."

Good responses: Those in which *clinical* symptoms such as fever, back pain, dysuria, frequency, urgency, etc., are relieved promptly and pyuria cleared.

Satisfactory responses: Those in which there was relief or alleviation of some of the presenting symptoms and a reduction but no complete clearing of pyuria.

Poor responses: Those in which there was no significant effect on the symptoms and no appreciable change in pyuria.

Science for the world's well-being[®] (Pfizer logo) Since 1849 Pfizer Laboratories Division, Chas. Pfizer & Co., Inc. New York, New York 10017 p159X67 (c) 1967, Chas. Pfizer & Co., Inc. Printed in U.S.A. Issued -----