

VIBRAMYCIN HAS BEEN FOUND CLINICALLY EFFECTIVE IN THE TREATMENT OF A VARIETY OF INFECTIONS CAUSED BY SUSCEPTIBLE STRAINS OF GRAM-POSITIVE AND GRAM-NEGATIVE BACTERIA. *

Site of Infection	Indications	Pathogens (Susceptible Strains)
Lower Respiratory Tract	- Streptococcus pneumoniae - Mycoplasma pneumoniae - Haemophilus influenzae - Brucella	- Staphylococcus aureus - Streptococcus E. coli - Klebsiella pneumoniae
Soft Tissue	- Impetigo - Furunculosis - Cellulitis - Abscess - Infected wounds - Paronychia	- Staph. aureus - Staph. albus - Streptococcus E. coli - Klebsiella-Aerobacter group
Genitourinary Tract	- Pyelonephritis - Cystitis - Urethritis	- Klebsiella-Aerobacter group E. coli - Enterococcus - Staphylococcus - Streptococcus - Neisseria gonorrhoeae

Since Vibramycin (doxycycline) is a member of the tetracycline series of antibiotics, it may be expected to be useful in the treatment of infections which respond to other tetracyclines. These include the following infections when caused by susceptible organisms:

Site of Infection	Indications	Pathogens (Susceptible Strains)
	Gastrointestinal Infections	E. histolytica - Shigella - Salmonella - Paratyphoid E. coli

ACNE VULGARIS AND ACNE CONGLOBATA

*In Combination with Streptomycin

*BECAUSE NOT ALL STRAINS OF THE LISTED PATHOGENS ARE SUSCEPTIBLE, IT IS RECOMMENDED THAT ROUTINE CULTURE AND SUSCEPTIBILITY STUDIES BE PERFORMED.