

Vibramycin doxycycline

an efficient oral broad-spectrum antibiotic
in terms of...

Serum concentrations of Vibramycin peak
at a rate which approaches that of a tetracycline
I.M. injection, indicating the great absorption of Vibramycin
from the G.I. tract.

Long half-life and slow urinary clearance of
Vibramycin allow you to prescribe it on a
one-dose-a-day basis after the first day.

FOR MORE SEVERE INFECTIONS, 100 MG. EVERY 12 HOURS IS RECOMMENDED.

The lowest daily dose of any oral tetracycline.

MINIMAL DOSE RELATED LOWER G.I. SIDE EFFECTS OBSERVED THUS FAR.

May be administered with meals or milk
without significant loss of activity.

ANTIMICROBIAL SPECTRUM WHICH IS COMPARABLE TO OTHER TETRACYCLINES.

References: 1. Research data on file, Pfizer Medical Department, Pfizer Laboratories. 2. English, A. R. and Lynch, J. E., Proc. Soc. Exp. Biol. Med. 124:586, Feb., 1967. 3. Clinical data on file, Pfizer Medical Department, Pfizer Laboratories. Available to physicians on request. 4. Rosenblatt, J. E., Barrett, J. E., Brodie, J. I. and Kirby, W. M. M.: Antimicrobial Agents and Chemotherapy, 1966, Ann Arbor, American Society for Microbiology, 1967, pp. 134-141. 5. Kunin, C. M., Dornbush, A. C. and Finland, M., J. Clin. Invest. 38:1950, Nov., 1959. 6. Migliardi, J. R. and Schoch von Wiltenau, M., presented at Int. Cong. Chemother., Vienna, June 26-July 1, 1967.

In dosage and absorption
Developed for Efficiency

