

Steri-Vial[®] No. 57

CHLOROMYCETIN[®]
(CHLORAMPHENICOL)
SODIUM SUCCINATE

FOR INTRAVENOUS ADMINISTRATION
Equivalent to **1 GRAM** chloramphenicol
CAUTION—Federal law prohibits dispensing without prescription.

PARKE-DAVIS
Stock 35-57-1

WARNING—Blood dyscrasias may be associated with the use of chloramphenicol. It is essential that adequate blood studies be made. See enclosed warnings and precautions.

Usual Adult Daily Dose
50 mg. per Kg. See package insert.

To prepare a 10% solution, add 11 cc. of sterile aqueous diluent such as Water for Injection or 5% Dextrose Injection. Solution should be used within 24 hours. A cloudy solution should not be used.

WARNING—Keep out of the reach of children.

Not suitable for laboratory diagnostic use.
To prepare a 10% solution, add 11 cc. of sterile aqueous diluent such as Water for Injection or 5% Dextrose Injection.
Usual Adult Daily Dose—50 mg. per Kg. See package insert.

CAUTION—Federal law prohibits dispensing without prescription.

WARNING—Blood dyscrasias may be associated with the use of chloramphenicol. It is essential that adequate blood studies be made. See enclosed warnings and precautions.

Parke, Davis & Company
Detroit, Michigan 48232 U.S.A.

Lot

Steri-Vial[®] No. 57

CHLOROMYCETIN[®]
(CHLORAMPHENICOL)
SODIUM SUCCINATE
FOR INTRAVENOUS ADMINISTRATION
1 GRAM

Stock 35-57-1

S-V No. 57

CHLOROMYCETIN[®]
(CHLORAMPHENICOL)
SODIUM SUCCINATE

FOR INTRAVENOUS ADMINISTRATION
EQUIVALENT TO 1 GRAM
CHLORAMPHENICOL

PARKE-DAVIS
Stock 35-57-1

Steri-Vial[®] No. 57

CHLOROMYCETIN[®]
(CHLORAMPHENICOL)
SODIUM SUCCINATE
FOR INTRAVENOUS ADMINISTRATION
1 GRAM

Expiration Date and Lot

WARNING—Keep out of the reach of children.

Patient:

Room:

Sterile solution may be kept at room temperature for 30 days without significant loss of potency. A cloudy solution should not be used.