TABLE 7.—RETROSPECTIVE STUDY OF THE RELATIONSHIP OF RHEUMATIC SYMPTOMS TO THE USE OF ORAL CONTRACEPTIVES (OC) IN 86 PATIENTS

45 PATIENTS USING OC AT TIME OF INITIAL SURVEY, SYMPTOMS PRECEDED OC USE (22)

	Symptom o			
	Increase	Equal	Decrease	Total
Rheumatoid arthritis	5	2	1	8
SLE	0	1	1	2
Arthralgia-myalgiaSynovitis	0	2	. 0	2
Miscellaneous	i	3	ĭ	5
 Total	9	10	3	22

## SYMPTOMS BEGAN DURING OC USE (23)

	Symptom cha	nge after st	0		
-	Increase	Equal	Decrease	Continued OC	Total
Rheumatoid arthritis	0 0 0 0	4 0 1 0	3 1 3 1	3 1 1 3	10 2 5 4
Miscellaneous	Ŏ	Ō	Ō	2	ż
Total	0	5	8	10	23

## 41 PATIENTS USED OC AT SOME TIME PRIOR TO SURVEY, SYMPTOMS PRECEDED USE OF OC (20)

	Temporal relationship symptoms (Sx) to OC use							
•	Sx change with use			Sx change after stopping				
	Increase	Equal	Decrease	Increase	Equal	Decrease	None 1	Total
Rheumatoid arthritis	1 3 0 1	6 0 3 3	0 0 0 0	0 0 0 0	6 1 3 3	1 2 0 1	1 0 2 0	8 3 5 5
Total	5	12	0	0	13	4	3	21

## SYMPTOMS FOLLOWED OC USE (21)

	Temporal relationship symptoms (Sx) to OC use					
_	Sx chan	ge after stop	ping		At . 1 . 6	Total
	Increase	Equal	Decrease	None 1	No infor- mation	
Rheumatoid arthritis	0	4	1	2	1	8
Arthralgia-myalgiaSynovitis	Ŏ 0	2	i 1	4 1	i i	8
Miscellaneous	0	0	0	1	Ö	1
Total	0	. 6	4	8	3	21

<sup>1</sup> No relationship, i.e., OC used and stopped more than 30 days preceding development of symptoms.

Senator McIntyre. The next witness is Dr. Edmond Kassouf, a practicing a physician from New Japan

ticing physician from New Jersey.

We are delighted to welcome you here. Please proceed in any manner you would like. Anything you can do in the way of paraphrasing or summarizing, we would appreciate it, but I want you to feel that the committee will hear you out.