

# Just what the doctor ordered

and spaced just right in the family plan ... worked out years before  
by the physician and the baby's parents. With the Enovid-E record  
of dependability (more than 4 million woman-years of actual use),  
this kind of planning becomes possible.

**Adverse Reactions**—A statistically significant association has been shown between use of oral contraceptives and the following serious adverse reactions: thrombophlebitis, pulmonary embolism.

Although available evidence is suggestive of an association, such a relationship has been neither confirmed nor refuted for the following serious adverse reactions: cerebrovascular accidents, neuro-ocular lesions, e.g., retinal thrombosis and optic neuritis.

The following adverse reactions are known to occur in patients receiving oral contraceptives: nausea, vomiting, gastrointestinal symptoms (such as abdominal cramps and bloating), breakthrough bleeding, spotting, change in men-

strual flow, amenorrhea during and after treatment, edema, chloasma or melasma, breast changes (tenderness, enlargement, secretion), change in weight, changes in cervical erosion and cervical secretions, suppression of lactation when given immediately post partum, cholestatic jaundice, migraine, allergic rash, rise in blood pressure in susceptible individuals, mental depression.

Although the following adverse reactions have been reported in users of oral contraceptives, an association has been neither confirmed nor refuted: anovulation post treatment, premenstrual-like syndrome, changes in libido, changes in appetite, cystitis-like syndrome, headache, nervousness, dizziness, fatigue, backache, hirsutism, loss of

scalp hair, erythema multiforme and nodosum, hemorrhagic eruption, itching.

The following laboratory results may be altered by oral contraceptives: hepatic function: increased sulobromophthalein and other tests; coagulation tests: increase in prothrombin, Factors VII, VIII, IX and X; thyroid function: increase in PBI and butanol extractable protein bound iodine, and decrease in T<sub>3</sub> uptake values; metyrapone test; pregnanediol determination.

References: 1. Inman, W. H. W., and Vessey, M. P.: Brit. Med. J. 2:193-199 (April 27) 1968. 2. Vessey, M. P., and Doll, R.: Brit. Med. J. 2:199-205 (April 27) 1968.

Before prescribing see complete prescribing information.

**makes planning life a little easier.**

**SEARLE** G. D. Searle & Co.  
P.O. Box 5110, Chicago, Illinois 60680

# Enovid-E®

norethynodrel 2.5 mg., mestranol 0.1 mg.