

TABLE II.—*Central nervous system symptoms*

Complications:	Cases
Severe vertigo:	2
With hospitalization for 1 week:	
(Enovid, 5 mg., for 5 months)-----	1
(Enovid, 5 mg., for 1 cycle)-----	1
Severe headache (Orthonovum, 2 mg., for 4 months)-----	1
Severe nausea and malaise (C-Quens for 10 days)-----	1
Total-----	4

TABLE III.—*Altered menstrual cycle after omission of pills*

Complications:	Cases
Metrorrhagia for 5 months (Enovid for 29 months)-----	1
Metrorrhagia for 24 months (Enovid-E for 6 months)-----	1
Oligo-amenorrhea for 3 months (Enovid for 9 months)-----	1
Amenorrhea for 6 months (Enovid for 3 years and then C-Quens for 6 months)-----	1
Total-----	4

TABLE IV.—*Infections*

Complications:	Cases
Vaginal:	
Monilial vulvovaginitis resistant to treatment until pills omitted (Enovid, 8 pts.; others, 3 pts.)-----	11
Membranous vaginitis (C-Quens for 4 months)-----	1
Nonspecific vaginitis (Enovid for 3½ years)-----	1
Systemic: Furunculosis, severe (Enovid for 14 months)-----	1
Total-----	14

TABLE V.—*Abnormal changes in pelvic organs*

Complications:	Cases
Pseudodecidual reaction of cervix (Enovid 8 pts., Norlestrin 1 pt.)--	19
Enlarging leiomyomas of uterus-----	2
(Enovid, 5 mg., for 6 months)-----	1
(Orthonovum, 2 mg. for 6 months)-----	1
Multiple endocervical polyps, pathology report suggested "as a result of Enovid" (Enovid, 5 mg., for 18 months)-----	1
Repeated episodes of cystitis (Enovid for 3½ years)-----	1
Total-----	13

TABLE VI.—*Vascular accidents*

Complications:	Cases
2 free appendices epiploicas found at surgery (Enovid for 6 months)---	1
3 pulmonary emboli (Orthonovum, 2 mg., for 2 months)-----	1
Thrombosis of vein of vulva (Oracon for 18 months)-----	1
Subarachnoid hemorrhage (Enovid-E for 3 months)-----	1
Total-----	4

¹ This change was seen in many additional patients not included in this table.