

- 38

Medical examiner case
0—No
1—Yes
2—Unknown
- 39

Oral contraceptive death
0—No
1—Yes
2—Not sure
- 40

On drug at time of death
0—Yes
1—None for 1 month
2—None for 2 months
3—None for 3 months
- 40

4—None for 4 months
5—None for 5 months
6—None for 6 months
7—None for 7 months
8—None for 8 months
9—None for 9 months or more
x—None for less than 1 month
y—Unknown
- 41

Indication for drug use
0—Contraception
1—Pregnancy test
- 41

2—Dysmenorrhea
3—Menorrhagia
4—Endometriosis
5—Irregular menses
6—Premenstrual tension
7—Habitual abortion
8—Ulcerative colitis
y—Unknown
- 42

Obese
0—No
1—Yes
2—Border line
y—Unknown

Cause	Less than 1	1-4	5-8	9-12	13-18	19-24	25-36	37-48	50	60	Un-known	Total
None listed	1	3	1	1		3	1		1			11
Pulmonary embolus	7	38	17	2	3						8	75
Cerebral embolization		1										1
Cerebral hemorrhage		3	1								3	7
Heart disease	1	1				1		1		1	1	6
Thrombocytopenia				1	1						1	3
General thrombosis	2											2
Leukemia											1	1
Peritonitis							1					1
Unknown		3										3

Race

	All		T.E.	
	Number	Percent	Number	Percent
White	66	60.0	43	66.2
Nonwhite	17	15.5	6	9.2
Unknown	27	24.5	16	24.6
Total	110		65	

Age

	All		T.E.	
	Number	Percent	Number	Percent
15-19	3	2.7	2	3.1
20-24	34	30.9	24	36.8
25-29	19	17.3	10	15.4
30-34	19	17.3	12	18.5
35-39	19	17.3	9	13.9
40-44	7	6.4	4	6.2
45 and over	4	3.6	3	4.6
Unknown	5	4.5	1	1.5
Total	110		65	
Range	18-50		18-50	
Average	30.2		28.5	