TABLE IV.—PREVALENCE RATES FOR CERVICAL CARCINOMA IN SITU COMPARED ACCORDING TO AGE AT FIRST PREGNANCY AND NUMBER OF LIVE BIRTHS IN WOMEN KNOWN TO BE USING THE DIAPHRAGM OR CONTRACEPTIVE ORAL STEROIDS FOR ONE YEAR OR LONGER

	Never pregnant	Age at first pregnancy			
		<19 years		20 years+	
Diaphragm: Number of cases Ca. in situ Total number of cases Prevalence rates per 1,000	_ 208		7 1, 106 6-3		2, 560 3.5
		Number of live births			
		0-2	3+	0–2	. 3+
Number of cases Ca. in situ Total number of cases Prevalence rates per 1,000		3 357 8-4	4 749 5⋅3	1, 365 2.2	1, 195 5.0
		Age at first pregnancy			
		<19 years		20 years+	
Oral steroids: Number of cases Ca. in situ Total number of cases Prevalence rates per 1,000	_ 414		29 2, 991 9.7		2, 926 11
		Number of live births			
		0–2	3+	0–2	3+
Number of cases Ca. in situ		1, 033 5.8	1, 958 12-0	16 1,595 10.0	1, 331 12-0

TABLE V.—PREVALENCE RATES FOR CERVICAL CARCINOMA IN SITU COMPARED ACCORDING TO NUMBER OF LIVE BIRTHS IN WOMEN KNOWN TO BE USING THE DIAPHRAGM OR CONTRACEPTIVE ORAL STEROIDS FOR ONE YEAR OR LONGER

	Number of live births		
	0-2	3+	
Diaphragm:			
Number of cases Ca. in situ	6	10	
Total number of cases	1,722	1,944	
Prevalence rates per 1,000Oral steroids:	3.3	2-1	
Number of cases Ca. in situ	22	39	
Total number of cases	2, 628	3, 289	
Prevalence rates per 1,000	8.4	12.0	

TABLE VI.—PREVALENCE RATES FOR CERVICAL CARCINOMAS IN SITU COMPARED ACCORDING TO ECONOMIC STATUS IN WOMEN KNOWN TO BE USING THE DIAPHRAGM OR CONTRACEPTIVE ORAL STEROIDS FOR ONE YEAR OR LONGER

	Family net income		
	<\$75/week	>\$75/week	
Diaphragm:			
Number of cases Ca, in situ	11	. 5	
Total number of cases	1,441	2, 433	
Prevalence rates per 1,000	7.6	2.1	
Oral steroids:			
Number of cases Ca. in situ	34	28	
Total number of cases	3, 310 10	28 3, 021	
Prevalence rates per 1,000	´ 10	′ 9.3	